

Draft Guidance on Betamethasone Dipropionate

This draft guidance, when finalized, will represent the current thinking of the Food and Drug Administration (FDA, or the Agency) on this topic. It does not establish any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the Office of Generic Drugs.

Active Ingredient: Betamethasone dipropionate

Dosage Form; Route: Lotion; topical

Recommended Studies: Two studies

1. Type of study: Pilot Vasoconstrictor Study
Design: A pilot dose duration-response study using the reference product under un-occluded conditions
Strength: EQ 0.05% Base
Subjects: Males and non-pregnant, non-lactating females, general population
Additional comments: Refer to the guidance “Topical Dermatological Corticosteroids: In Vivo Bioequivalence”.

-
2. Type of study: Pivotal vasoconstrictor study
Design: A pivotal bioequivalence study under un-occluded conditions
Strength: EQ 0.05% Base
Subjects: Males and non-pregnant, non-lactating females, general population
Additional Comments: See comments above
-

Analytes to measure (in appropriate biological fluid): Not applicable

Bioequivalence based on (90% CI): Pivotal vasoconstrictor study

Waiver request of in vivo testing: Not applicable

Dissolution test method and sample times: Not applicable