

Contains Nonbinding Recommendations

Draft – Not for Implementation

Draft Guidance on Nateglinide

October 2024

This draft guidance, when finalized, will represent the current thinking of the Food and Drug Administration (FDA, or the Agency) on this topic. It does not establish any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the Office of Generic Drugs.

In general, FDA’s guidance documents do not establish legally enforceable responsibilities. Instead, guidances describe the Agency’s current thinking on a topic and should be viewed only as recommendations, unless specific regulatory or statutory requirements are cited. The use of the word *should* in Agency guidances means that something is suggested or recommended, but not required.

Active Ingredient: Nateglinide

Dosage Form: Tablet

Route: Oral

Strengths: 60 mg, 120 mg

Recommended Study: One in vivo bioequivalence study with pharmacokinetic endpoints

1. Type of study: Fasting
Design: Single-dose, two-treatment, two-period crossover in vivo
Strength: 120 mg
Subjects: Healthy males and non-pregnant, non-lactating females
Additional comments: All subjects should fast overnight for at least 10 hours prior to dosing and for 4 hours after dosing. A single oral dose (120 mg) should be administered with 240 mL of 20% glucose solution. Since, multiple plasma concentration peaks were often observed under fasting conditions, please ensure that the same sampling schedule is followed during the study for both test and reference drug administration. Females of reproductive potential should use effective contraception during the study.

Analyte to measure: Nateglinide in plasma

Bioequivalence based on (90% CI): Nateglinide

Waiver request of in-vivo testing: 60 mg strength based on (i) acceptable bioequivalence study on the 120 mg strength, (ii) acceptable in vitro dissolution testing of both strengths, and (iii) proportional similarity of the formulations between both strengths

Dissolution test method and sampling times: The dissolution information for this drug product can be found in the FDA’s Dissolution Methods database, <http://www.accessdata.fda.gov/scripts/cder/dissolution/>. Conduct comparative dissolution testing on 12 dosage units for each of both strengths of the test product and reference listed drug (RLD).¹ Specifications will be determined upon review of the abbreviated new drug application.

Document History: Recommended May 2007; Finalized May 2008; Revised October 2024

Unique Agency Identifier: PSG_021204

¹ If the RLD is not available, refer to the most recent version of the FDA guidance for industry on *Referencing Approved Drug Products in ANDA Submissions*.