



E14/S7B Implementation Working Group

E14/S7B IWG: Q&As: The Clinical Evaluation of QT/QTc Interval Prolongation and Proarrhythmic Potential for Non-Antiarrhythmic Drugs Questions and Answers

Current Version

dated 20 August 2026

In order to facilitate the implementation of the ICH E14/S7B IWG Guidelines, the ICH Experts have developed a series of Q&As:

E14/S7B Q&As Document History

Code	History	Date
ICH E14 Q&As	Approval by the ICH Steering Committee under <i>Step 4</i>	4 June 2008
ICH E14 Q&As (R1)	Approval by the ICH Steering Committee under <i>Step 4</i> of four newly added Questions on: Sex Differences; Incorporating New Technologies; Late Stage Monitoring; and Heart Rate Correction.	5 April 2012
ICH E14 Q&As (R2)	Approval by the ICH Steering Committee under <i>Step 4</i> of four newly added Questions on: Concentration-Response Relationships; Combination Products; Large Targeted Proteins and Monoclonal Antibodies; and Special Cases	21 March 2014
ICH E14 Q&As (R3)	Approval by the ICH Assembly under <i>Step 4</i> of the revision of the Question #5.1 on: <i>Use of Concentration Response Modeling of QTc Data</i>	10 December 2015
ICH E14/S7B Q&As	Approval by the ICH Assembly under <i>Step 4</i>	21 February 2022
ICH E14/S7B Q&As (R1)	Approval by the ICH Assembly under <i>Step 4</i> of the revision of E14 Questions 5.1, 6.1, and 6.3 and revised or introduced new Questions for Sections 1 and 3 for S7B.	20 August 2026

References

ICH E14	The Clinical Evaluation of QT/QTc Interval Prolongation and Proarrhythmic Potential for Non-Antiarrhythmic Drugs	May 2005
ICH E14	Questions & Answers (R3)	Dec 2015
ICH S7B	Nonclinical Evaluation of the Potential for Delayed Ventricular Repolarization (QT Interval Prolongation) by Human Pharmaceuticals	May 2005

Legal Notice: This document is protected by copyright and may be used, reproduced, incorporated into other works, adapted, modified, translated or distributed under a public license provided that ICH's copyright in the document is acknowledged at all times. In case of any adaption, modification or translation of the document, reasonable steps must be taken to clearly label, demarcate or otherwise identify that changes were made to or based on the original document. Any impression that the adaption, modification or translation of the original document is endorsed or sponsored by the ICH must be avoided. The document is provided "as is" without warranty of any kind. In no event shall the ICH or the authors of the original document be liable for any claim, damages or other liability arising from the use of the document. The above-mentioned permissions do not apply to content supplied by third parties. Therefore, for documents where the copyright vests in a third party, permission for reproduction must be obtained from this copyright holder.

Table of Contents

SECTION 1: E14 QUESTIONS AND ANSWERS	5
1. ELECTROCARDIOGRAMS METHODOLOGY	5
2. GENDER.....	11
3. POSITIVE CONTROL	12
4. STUDY DESIGN	14
5. USE OF CONCENTRATION RESPONSE MODELING OF QTC DATA.....	15
6. SPECIAL CASES	19
7. ELECTROCARDIOGRAMS MONITORING IN LATE STAGE CLINICAL TRIALS.....	22
SECTION 2: S7B QUESTIONS AND ANSWERS	25
1. INTEGRATED RISK ASSESSMENT.....	26
2. BEST PRACTICE CONSIDERATIONS FOR <i>IN VITRO</i> STUDIES	33
3. BEST PRACTICE CONSIDERATIONS FOR THE <i>IN VIVO</i> QT STUDIES.....	40
4. PRINCIPLES FOR PROARRHYTHMIA MODELS.....	44

SECTION 1: E14 QUESTIONS AND ANSWERS

1. ELECTROCARDIOGRAMS METHODOLOGY

#	Date of Approval	Questions	Answers
1.1	June 2008	Please discuss who should read electrocardiograms (ECGs), including the number and training of readers and the need for readers to be blinded.	The document recommends that the reader should be skilled but does not identify specific training that is needed. A technician reading with cardiologist over-read would certainly be consistent with the guidance. The attempt of the guidance to limit the number of readers represented an attempt to increase consistency. The guidance asks for assessment of intra- and inter-reader variability and suggests "a few skilled readers" (not necessarily a single reader) to analyse a whole thorough QT study, since many readers may increase variability. Training would be another way to improve consistency. It is recommended for the thorough QT Study that core ECG laboratories blind subject, time and treatment in order to reduce potential bias. The T wave analysis, which calls for all 12 leads, can be performed after the QT analyses, and requires comparison to the baseline ECG; it can, however, be blinded as to treatment.
1.2	June 2008	What is the position of ICH regarding the role of the following reading methods in the thorough QT/QTc study and other clinical trials? • fully manual • fully automated • manual adjudication (manual over-read, computer-assisted, semi-automated)	The techniques currently in use for the measurement of ECG intervals can be classified into three broad categories: fully manual, fully automated, and manual adjudication. Within each of these general categories, many different methodologies are subsumed that differ in terms of lead selection, the conventions used for defining T wave offset, and the criteria for the inclusion and exclusion of U waves. ECG readings can be performed on the following waveform presentations: • Raw waveforms: ECG waveforms recorded from a single lead;

			• Representative waveforms (median beats, reference cycles): Compositional waveforms constructed by a computer-based averaging process that involves aligning and combining data from all dominant, normally conducted raw ECG waveforms from a single lead; • Global Waveforms: Composite representation of cardiac electrical activity constructed by superimposing representative waveforms from all or several simultaneously recorded leads to form a spatial-vector complex, by weighted averaging of individual representative complexes with low noise and long duration, or by other methods. Fully Manual When using a fully manual reading technique, a human reader is responsible for examining the ECG waveform and placing the fiducial points to mark the beginning and the end of the intervals, without the assistance of a computer algorithm. Fully manual methods of fiducial point placement can be applied to raw, representative, and global waveforms. When fully manual measurements are made from the raw ECG waveforms in a single lead, three or more cycles should be averaged where available to produce the final determination of interval duration. An advantage of this approach is that the reader will not be influenced by prior computer placement of the fiducial points, but a weakness can be inter- and intra-reader variability, especially when measurements are performed over an extended time period (e.g., several months). Laboratories using manual reading techniques should observe standard operating procedures based on prospectively defined criteria for determining where the fiducial points should be placed. All readers in the laboratory should be trained in the consistent application of these criteria. Fully Automated Fully automated reading methods rely entirely upon a computer algorithm for the placement of the fiducial points and the measurement of the ECG intervals. Automated ECG interval measurements can be performed on raw,
--	--	--	--

		representative, or global ECG waveforms. Most digital electrocardiographs are equipped with algorithms that perform measurements on global waveforms. Although automated methods have the advantage of being consistent and reproducible, they can yield misleading results in the presence of noise or when dealing with abnormal ECG rhythms, low amplitude P or T waves, or overlapping U waves. The techniques used for construction and measurement of representative waveforms and global waveforms vary between different computerised algorithms and between different software versions within individual equipment manufacturers. As a result, between-algorithm and within-manufacturer variability of fully automated measurements can confound serial comparisons when the equipment or algorithm is not constant. Manual Adjudication (Manual Over-Read/Computer-Assisted/Semi-Automated) The manual adjudication approach refers to reading methods in which a computer algorithm is responsible for the initial placement of the fiducial points on the ECG waveform. A human reader subsequently reviews the algorithmic placement of the fiducial points, performing adjustments wherever the computerised measurements are considered to be inaccurate. This approach can have the advantage of greater consistency and reproducibility than fully manual readings, while providing an opportunity to correct any mistakes made by the algorithmic readings. Laboratories using manual adjudication techniques should observe standard operating procedures based on prospectively defined criteria for determining when fiducial points should be corrected. All readers in the laboratory should be trained in the consistent application of these criteria. The adjudication procedure should normally be performed on all waveforms being used for interval determination. If an alternative approach is used, such as adjudication limited to outlier intervals above and below a reference range, this methodology should be validated as described in Question 1.3.
--	--	---

			The ICH E14 Guideline currently recommends either fully manual or manual adjudication approaches for clinical trials in which the assessment of ECG safety is an important objective, such as the thorough QT/QTc study. When the thorough QT study is positive, fully manual or manual adjudication methods are currently recommended for an adequate sample of patients in late phase studies (see Section 2.3 in E14 document). When the thorough QT/QTc study is negative, routine ECG safety assessments in late phase clinical trials using fully automated reading methods would be adequate.
1.3	June 2008	The ICH E14 Guideline contains the following statement: “If well-characterised data validating the use of fully automated technologies become available, the recommendations in the guidance for the measurement of ECG intervals could be modified.” What would be expected of a sponsor that wished to validate and apply an automated reading method for regulatory submissions?	Efforts to develop more sophisticated and reliable methods for automated ECG readings for both QT interval and T wave morphology assessment are encouraged. There are at present no large scale studies to validate the use of fully automated reading methods in patients; however, there are examples of thorough QT/QTc studies in healthy volunteers in which automated methods have been used and validated for QT interval measurements against manual methods. QT Interval measurement There are at present no clear and widely accepted criteria for validation of new semi-automated or automated methods, but it is expected that each would be validated independently for its ability to detect the QT/QTc prolongation effects of drugs that are near the threshold of regulatory concern. Data supporting the validation of a new method should be submitted and could include descriptive statistics, Bland-Altman plots of agreement, superimposed plots of the baseline- and placebo- adjusted QTc and the RR as a function of time, together with data from any trials that have employed the method. T wave morphology assessment The suitability of automated ECG reading techniques for the assessment of morphological abnormalities has not yet been demonstrated. If a sponsor intends to develop a fully automated approach, without visual assessment for morphological changes, validation studies should include a demonstration

			that the automated method is capable of reading and interpreting a test set of abnormal ECGs correctly (e.g., abnormalities of T wave morphology, overlapping U waves). As with methods for QT interval determination, there are at present no clear and widely accepted criteria for validation of novel methods. Because changes in morphology can affect interval measurement, fully manual or manual adjudication (as defined in Question 1.2) techniques should be performed if treatment-emergent changes in morphology are observed. If, on the other hand, no morphology changes are observed, this would support the use of automated methodologies, provided they have been validated.
1.4	April 2012	How does a sponsor incorporate new technology or validate new methodology into the measurement and/or analysis of the QT interval?	The ICH process is better suited to the determination of regulatory policy once the science in a particular area has become more or less clear. In general, it is not well-suited to the qualification or validation of new technology. Sections 2.5.1 and 2.5.2 of the ICH E14 Guideline are rather discouraging about methodology outside conventional carts and human-determined measurements. Since ICH E14 was issued, 12-lead continuous recording devices have largely supplanted cart recorders in thorough QT studies without a formal validation process because of their performance in the context of a positive control. The impact of other innovative technologies can be assessed in studies incorporating a positive control. While some technologies could be assessed using other techniques in the absence of a positive control, this is more complex and beyond the scope of this Q&A. 12-lead continuous recording devices and other new technologies can be used in late phase clinical trials. Even though a positive control is not used in late stage studies, the new technology could be validated in other studies (such as the thorough QT study). In cases where a thorough QT study is not done, a sponsor can provide alternative methods for validating the technology.

1.5	April 2012	The ICH E14 Guideline states that QT interval corrected by Fridericia's and Bazett's correction should be submitted in all applications; is this still necessary? Is there a recommended approach to QT correction that is different than that specified in ICH E14?	Changes in heart rate could variably influence a drug's effect on repolarisation (i.e., QT interval) and correction methods with different characteristics are often applied. The principles set below would be applicable in all clinical studies (thorough QT or other studies). In adults, Bazett's correction has been clearly shown to be an inferior method of correcting for differences in heart rate among and within subjects. Therefore, QT interval data corrected using Bazett's corrections is no longer warranted in all applications unless there is a compelling reason for a comparison to historical Bazett's corrected QT data. Presentation of data with a Fridericia's correction is likely to be appropriate in most situations but other methods could be more appropriate. There is no single recommended alternative (see Q&A 1.4 on Incorporating New Technologies), but the following are some considerations. 1. Analyses of the same data using different models for correcting QT can generate discordant results. Therefore, it is important that the method(s) of correction, criteria for the selection of the method of correction, and rationale for the components of the method of correction be specified prior to analysis to limit bias. Model selection ought to be based on objective criteria and consider the uncertainty in parameter estimates. Alternative methods of correction should be used only if the primary method fails the pre-specified criteria for selection of the method of correction. 2. Corrections that are individualised to a subject's unique heart rate QT dynamic are not likely to work well when the data are sparse or when the baseline data upon which the correction is based do not cover at least the heart rate range observed on study drug.
-----	------------	---	---

2. GENDER

#	Date of Approval	Questions	Answers
2.1	June 2008	There are recognised differences in the baseline QTc between men and women. These were noted in early versions of the guidance. In E14, however, it is recommended that outliers be categorised as >450, >480 and >500 ms, regardless of gender. Can you say why there is no gender difference in the recommendation?	The 450, 480, and 500 ms categories refer to the values the E14 document suggests sponsors might use in characterising outliers. The numbers previously specified for males and females referred to “normal” QTc values, which may differ for men and women. This section was not included in the final document, however, and such considerations would be largely irrelevant to larger durations (e.g., 480, 500 ms). As the thorough QT/QTc study is designed to examine the propensity of a drug to prolong the QTc interval, it is appropriate to perform the study in male or female healthy volunteers.
2.2	April 2012	Should we enroll both sexes in a thorough QT study, and does the study need to be powered for independent conclusions about each sex?	Post-pubertal males have lower heart-rate corrected QT intervals than do pre-pubertal males or females generally. Women are generally smaller than men, so their exposure to a given fixed dose of a drug will generally be higher, and, if a drug prolongs QT, it can be expected to prolong it more in women because of the higher exposure. It is not settled whether and how often there are sex differences in response to QT-prolonging drugs not explained by exposure alone. The thorough QT study is primarily intended to act as a clinical pharmacology study in a healthy population using a conservative primary objective defining the drug’s effect on QT. It is unlikely that any of a variety of baseline demographic parameters would introduce a large difference in QT response to a drug in subpopulations defined by factors such as age, co-morbidity, and gender that is not explained by exposure. It is encouraged, but not mandatory, to include both men and women in the thorough QT study. Analyses of Concentration-Response Relationship (CRR) by sex can be helpful for studying the effect of the drug on QT/QTc interval in cases where there is evidence or mechanistic theory for a gender

			difference. However, the primary analysis of a thorough QT study should be powered and conducted on the pooled population. If the primary analysis is negative and if there is no other evidence suggesting gender differences, subgroup analysis by sex is not expected.
--	--	--	---

3. POSITIVE CONTROL

#	Date of Approval	Questions	Answers
3.1	June 2008	The ICH E14 Guideline emphasises the importance of assay sensitivity and recommends the use of a positive control. In order to accept a negative thorough QT/QTc study, assay sensitivity should be established in the study by use of a positive control with a known QT-prolonging effect. Please clarify how to assess the adequacy of the positive control in the thorough QT study.	The positive control in a study is used to test the study's ability (its "assay sensitivity") to detect the study endpoint of interest, in this case QT prolongation by about 5 ms. If the study is able to detect such QT prolongation by the control, then a finding of no QT effect of that size for the test drug will constitute evidence that the test drug does not in fact prolong the QT interval by the amount of regulatory concern. There are two conditions required for ensuring such assay sensitivity: 1. The positive control should show a significant increase in QTc; i.e., the lower bound of the one-sided 95% Confidence Interval (CI) must be above 0 ms. This shows that the trial is capable of detecting an increase in QTc, a conclusion that is essential to concluding that a negative finding for the test drug is meaningful; 2. The study should be able to detect an effect of about 5 ms (the QTc threshold of regulatory concern) if it is present. Therefore, the size of the effect of the positive control is of particular relevance. With this aim, there are at least two approaches: a. To use a positive control showing an effect of greater than 5 ms (i.e., lower bound of a one-sided 95% CI >5 ms). This approach has been proven to be useful in many regulatory cases. However, if the positive control has too large an effect, the study's ability to detect a 5 ms QTc prolongation

			might be questioned. In this situation, the effect of the positive control could be examined at times other than the peak effect to determine whether an effect close to the threshold of regulatory concern can be detected; b. To use a positive control with an effect close to 5 ms (point estimate of the maximum mean difference with placebo close to 5 ms, with a one-sided 95% CI lower bound >0). In using positive controls with smaller effects it would be very important to have a reasonably precise estimate of the drug's usual effect. Importantly, whatever approach is used, the effect of the positive control (magnitude of peak and time course) should be reasonably similar to its usual effect. Data suggesting an underestimation of QTc might question the assay sensitivity, thus jeopardising the interpretability of the thorough QT study results.
3.2	June 2008	Please clarify the need for blinding the positive control in the thorough QT study.	The use of a double-blinded positive control does not appear to be essential, provided that the reading of ECGs is performed in a blinded manner as described in Question 1.2 and the study is carefully designed to ensure that specified study procedures are followed uniformly. This means that the same protocol for administering the test drug and placebo, taking blood samples and collecting the ECG data should also be used when giving the positive control. This does not mean that other aspects of the study, such as the duration of treatment with the positive control and the other treatment groups, would be identical. If blinding of the positive control is performed, common methods include the use of double-dummy techniques and over-encapsulation.

4. STUDY DESIGN

#	Date of Approval	Questions	Answers
4.1	June 2008	In ICH E14, the recommended metric to analyse for a cross-over study is the largest time-matched mean difference between the drug and placebo (baseline-adjusted) over the collection period. Please discuss the most appropriate metric to assess a drug's effect on QT/QTc interval when the data are collected in a placebo-controlled parallel design study (i.e., when there is no corresponding placebo value for each patient).	Regardless of the study design, “the largest time-matched mean difference between drug and placebo (baseline-adjusted)” is determined as follows: The mean QTc for the drug (i.e., averaged across the study population) is compared to the mean QTc for placebo (averaged across the study population) at each time point. The “largest time-matched mean difference between drug and placebo” is the largest of these differences at any time point. The term “baseline-adjusted” in ICH E14 implies that the baseline data are taken into account in the statistical analysis. Differences in baseline assessment between cross-over and parallel design studies are discussed in Question 4.2.
4.2	June 2008	Please discuss the need for baseline measurements, and when needed, how they should be collected, for cross-over and parallel design thorough QT studies.	Adjustment for baseline measurements is potentially useful for several purposes, including detection of carry-over effects, reducing the influence of inter-subject differences and accounting for diurnal effects such as those due to food. There is no single best approach for baseline adjustment, but all planned baseline computations should be prospectively defined in the clinical trial protocol. Two kinds of baseline are commonly used: “time-matched” baseline (taken at exactly the same time-points on the day prior to the beginning of treatment as on the treatment day) and “pre-dose” baseline (taken shortly prior to dosing). The “pre-dose” baseline is used for adjustment for inter-subject differences but not for diurnal effects. The choice of baseline is influenced by whether the study is parallel or cross-over. For a parallel-group study a time-matched baseline allows the detection of differences in diurnal patterns between subjects that would not be detected

			by a predose baseline. In a parallel study a “time-matched” baseline day, if performed, would ideally occur on the day before the start of the study. In contrast, in a cross-over study a time-matched baseline is usually not necessary because adjustments for subject- and study-specific diurnal variation are implicit by design in the assessment of time-matched drug-placebo differences in QT/QTc effect. The “pre-dose” baseline is therefore usually adequate for cross-over studies. Obtaining replicate ECG measurements (for example, the average of the parameters from about 3 ECGs) within several minutes of each nominal time point at baseline and at subsequent times will increase the precision of the estimated changes in QT/QTc effect.
--	--	--	---

5. USE OF CONCENTRATION RESPONSE MODELING OF QTC DATA

#	Date of Approval	Questions	Answers
5.1	June 2026	The ICH E14 Guideline states (in Section 3.2.3, page 10) that analysis of the relationship between drug concentration and QT/QTc interval changes is under active investigation. Has this investigation yielded a reasonable approach to concentration-response modeling during drug development? How can assessment of the concentration-response relationship guide the interpretation of QTc data?	Concentration-response analysis, in which all relevant data across all doses are used to characterise the potential for a drug to influence QTc, can serve as an alternative to the by-time-point analysis or intersection-union test as the primary basis for decisions to classify the risk of a drug. In either case this result is an important component of the totality of evidence assessment of the risk of QT prolongation. The overall assessment of risk of QT prolongation includes nonclinical data, the time course of QT prolongation, the magnitude of QT prolongation, categorical analyses of outliers, and certain adverse events in patients that can signal potential proarrhythmic effects. There are many different types of models for the analysis of concentration-response data, including descriptive pharmacodynamic (PD) models (e.g., linear or Emax models), or empirical models that link pharmacokinetic (PK) models (dose-concentration-response) with PD

		models. It is recognised that concentration-response analyses of the same data using models with different underlying assumptions can generate discordant results. Therefore, it is important that the modeling methods and assumptions, criteria for model selection, rationale for model components, and potential for pooling of data across studies be specified prior to analysis to limit bias. Prospective specification of model characteristics (e.g., structural model, objective criteria, goodness of fit) based on knowledge of the pharmacology is recommended whenever possible. On occasion, the QT effect is not a direct function of plasma concentration. For example, drugs that cause QT prolongation as a result of changes in protein synthesis or trafficking or drugs with accumulation into myocardial tissues might demonstrate hysteresis. Testing for model assumptions, hysteresis (a plot of data by-time point and a hysteresis loop plot), and goodness of fit should be documented. Concentration-response analysis can be challenging when more than one molecular entity-multiple drugs or parent plus metabolites-contributes to the QTc effect. Dose and exposure definitions • Therapeutic dose: dose evaluated in Phase 3 trial or recommended in product labeling • Clinical exposure: mean steady state maximum concentration ($C_{\max,ss}$) associated with the maximum therapeutic dose • High clinical exposure: exposure ($C_{\max,ss}$) achieved when the maximum therapeutic dose is administered in the presence of the intrinsic or extrinsic factor (e.g. organ impairment, drug-drug interaction, food effect, etc.) that has the largest effect on increasing $C_{\max,ss}$ • Supratherapeutic dose: dose that provides exposures (mean $C_{\max}$) exceeding the high clinical scenario Important considerations
--	--	---

			Concentration-response data need not come from a dedicated QT study, nor even a single study, but there are several new and important considerations. 1. Data can be acquired from first-in-human (FIH) studies, multiple-ascending dose (MAD) studies, or other studies provided that the concentrations that can be safely achieved are well above the exposure at the maximum therapeutic dose at steady-state, and reflect high clinical exposure scenario situations such as drug-drug and drug-food interactions, organ dysfunction, and/or genetically impaired metabolism. Whenever possible, sponsors are encouraged to explore a wide dose range in early phase studies to enable characterisation of effects at high concentrations. 2. Efficient concentration-response analysis using data acquired in studies with other purposes requires as much quality control as is needed for a dedicated study. This includes robust, high-quality electrocardiogram (ECG) recording and analysis sufficient to support a valid assay for ECG intervals (see ICH E14 and Q&A #1). 3. If there is an intention to pool ECG interval data from multiple studies, it is important to test for heterogeneity. Pooling of studies that were not planned for this purpose can produce bias. This potential should be critically discussed in the analysis plan. 4. A separate positive control would not be necessary if either of the following conditions is met: a. There are data characterising the response at a sufficient multiple (commonly 2x) of the high clinical exposure (see ICH E14 Section 2.2.2); b. If the high clinical exposure has been achieved in the clinical ECG assessment, but a sufficient multiple has not been obtained (e.g., for reasons of safety or tolerability, saturating absorption, etc.), then a nonclinical integrated risk assessment can be used as supplementary evidence. The reason higher doses were not tested should be adequately justified. The nonclinical integrated risk
--	--	--	--

			assessment should follow the testing strategy defined in ICH S7B Q&A 1.4, parts b and c and show low risk as defined in ICH S7B Q&A 1.2. Decision-making Both the intersection-union test and the concentration-response analysis can estimate the maximum effect of a drug treatment on the QTc interval, but they are not used to test the same hypothesis. As mentioned above, inspection of the time course of QT prolongation is important. However, hypothesis testing based on a by-time point analysis (intersection-union test or point estimate and confidence intervals) is inappropriate in studies designed for a concentration-response analysis, if not powered to assess the magnitude of QT prolongation for each time point. When using a concentration-response analysis as the primary basis for decisions to classify the risk of a drug, the upper bound of the two-sided 90% confidence interval for the QTc effect of a drug treatment as estimated by exposure-response analysis should be <10 ms at the highest clinically relevant exposure to conclude that an expanded ECG safety evaluation during later stages of drug development is not needed. (See ICH E14, Section 2.2.4 and Q&A #7). Other uses In addition to serving as the basis for regulatory decision-making, concentration-response analysis has established its utility in several settings enumerated below. <u>Providing insight into regimens not studied directly</u> An understanding of the concentration-response relationship can help predict the QT effects of doses, dosing regimens, routes of administration, or formulations that were not studied directly. Interpolation within the range of concentrations studied is more reliable than extrapolation above the range. <u>Predicting QTc effects of intrinsic and extrinsic factors that affect PK</u>
--	--	--	--

			Understanding the concentration-response relationship can help predict the effects of intrinsic (e.g., cytochrome P450 isoenzyme status) or extrinsic (e.g., drug-drug PK interactions) factors, possibly affecting inclusion criteria or dosing adjustments in later phase studies.
--	--	--	--

6. SPECIAL CASES

#	Date of Approval	Questions	Answers
6.1	June 2026	The ICH E14 Guideline states that in certain cases a conventional thorough QT study might not be feasible. In such cases what other methods should be used for evaluation of QT/QTc and proarrhythmic potential?	An integrated nonclinical and clinical QT/QTc risk assessment can be particularly valuable when a thorough QT study or concentration-QTc analysis meeting similar quality control as needed for a dedicated study as described in 5.1 is not feasible. This situation can arise under scenarios where a placebo-controlled comparison is not possible; safety considerations preclude administering supratherapeutic doses to obtain high clinical exposures and/or safety or tolerability prohibit the use of the product in healthy participants. The design elements that include placebo and healthy participant dosing assist in decreasing variability, but their absence does not preclude interpretation. The integrated nonclinical and clinical QT/QTc risk assessment should include: 1. Nonclinical data according to the testing strategy defined in ICH S7B Q&A 1.4, parts b and c and any follow-up nonclinical studies, especially those selected to overcome the challenges encountered in the clinical studies (see ICH S7B Q&A 1.2); and

			2. Alternative QT clinical study designs incorporating ECG assessments with as many of the usual “thorough QT/QTc” design features as possible (see ICH E14 Section 2.2 and Q&A 5.1). An integrated QT/QTc risk assessment can also be particularly valuable for drugs with confounding heart rate effects that could impact accurate determination of the QTc. Advanced methodologies for controlling (e.g., pacing) or correcting for heart rate changes in the nonclinical studies and/or conducting QTc assessments in the intended patient population might be informative in this situation. If tolerance to the chronotropic effect develops with repeat dosing, upward titration regimens can sometimes be employed to avoid or minimise the confounding effects of drug-induced heart rate changes on the QTc assessment. Decision-Making A totality of evidence argument based on the results of an integrated nonclinical and clinical QT/QTc assessment could be made at the time of marketing application. A drug that meets the following criteria would be considered to have a low likelihood of proarrhythmic effects due to delayed repolarisation: 1. The non-clinical data obtained according to the testing strategy in ICH S7B Q&A 1.4, parts b and c, show low risk as defined in ICH S7B Q&A 1.2. 2. The high-quality ECG data (see ICH E14 and ICH E14 Q&A 1) collected in the alternative QT clinical assessment do not suggest QT prolongation, generally defined under this Q&A as an upper bound of the two-sided 90% confidence interval around the estimated maximal effect on ΔQT_c less than 10 ms, as computed by the concentration-response analysis or the intersection-union test. If applicable, there should be no notable imbalances between treatment/dose arms in the proportion of subjects exceeding outlier thresholds.
--	--	--	--

			3. A cardiovascular safety database that does not suggest increased rate of adverse events that signal potential for proarrhythmic effects (ICH E14 Section 4). When justified, a totality of evidence argument for a drug to have a low likelihood of proarrhythmic effects due to delayed repolarisation could still be made for a drug that has an upper bound of the two-sided 90% confidence interval around the estimated maximal effect on ΔQ_{Tc} of 10 ms or more. The determination will depend on the quality and details of the clinical data (e.g., estimated ΔQ_{Tc} mean and upper bound values, slope of any concentration-ΔQ_{Tc} relationship) and nonclinical data (e.g., difference between the hERG safety margin, as applicable, for the investigational drug and the threshold for defining low risk). If nonclinical studies do not show low risk (or are not performed), there is reluctance to conclude a lack of an effect in an absence of a positive control; however, if the upper bound of the two-sided 90% confidence interval around the estimated maximal effect on ΔQ_{Tc} is less than 10 ms, the treatment is unlikely to have an actual mean effect as large as 20 ms.
6.2	March 2014	The ICH E14 Guideline does not address the approach to QT measurement during drug development in the case of combination drug products. Is it recommended that measurement of QT prolongation be performed on drug combinations?	In general, combinations of two or more drugs are unlikely to need a thorough QT/Q_{Tc} study or intensive late stage monitoring, if the component drugs have been demonstrated to lack relevant effects in thorough QT/Q_{Tc} studies as described in ICH E14. If one or more of the component drugs have not been individually characterised for effects on the QT/Q_{Tc} interval, they may be evaluated in combination or independently.

6.3	June 2026	Are sponsors expected to conduct thorough QT studies as part of the development of therapeutic modalities (non-small molecules) where the potential to induce delayed ventricular repolarisation via interaction with the hERG channel blocking is not likely?	For certain modalities, including monoclonal antibodies and large targeted proteins as described in ICH S7B Q&A 1.4 part a, the potential to induce delayed ventricular repolarisation via interaction with the hERG channel is not likely. In these circumstances, a thorough QT/QTc study is not necessary, unless the potential for proarrhythmic risk is suggested by mechanistic considerations or data from clinical or non-clinical studies.
------------	------------------	--	---

7. ELECTROCARDIOGRAMS MONITORING IN LATE STAGE CLINICAL TRIALS

#	Date of Approval	Questions	Answers
7.1	April 2012	The ICH E14 Guideline describes in Section 2.3 (Clinical Trial Evaluation After the “Thorough QT/QTc Study”) that “adequate ECG assessment to accomplish this [monitoring] is not fully established.” Is there now a reasonable approach to evaluating QTc in late stage clinical development in the case of a finding of QT prolongation prior to late phase studies?	Clarification of Approach to Evaluating QTc in Late Stage Clinical Development The purpose of a thorough QT study is to characterise the effect of the drug on ventricular repolarisation (QT interval). It is not the purpose of the thorough QT study to assess the risk of Torsade de Pointes (TdP) in the target population, but rather to determine whether further data are warranted to assess risk. A finding of QT prolongation above the regulatory threshold of interest (a positive thorough QT study) might call for further electrocardiographic follow up in late phase studies. The extent of the follow up would be affected by the magnitude of the estimated prolongation at doses and concentrations at which this occurs. If prolongation is substantial at concentrations expected to occur in clinical studies, it is important to protect patients in later trials and to obtain further information on the frequency of marked QT prolongation. In some cases in which there is a large margin of safety between therapeutic exposures and the exposures that

			result in significant ECG interval changes, an intensive ECG follow-up strategy might not be warranted. The recommended intensity of the monitoring and assessment in late-stage trials will depend on: A. The magnitude of QTc prolongation seen in the thorough QT study or early clinical studies. B. The circumstances in which substantial QT prolongation might occur (i.e., in ordinary use or only when drug concentrations are markedly increased [e.g., by renal or hepatic impairment, concomitant medications]). C. Pharmacokinetic properties of the drug (e.g., high inter-individual variability in plasma concentrations, metabolites). D. Characteristics of the target patient population that would increase the proarrhythmic risk (e.g., structural heart disease). E. The presence of adverse effects that can increase proarrhythmic risk (e.g., hypokalemia, bradycardia, heart block). F. Other characteristics of the drug (e.g., pharmacodynamics, safety pharmacology, toxicology, drug class, hysteresis). The following examples delineate the scope of recommended ECG investigations based on outcome of the thorough QT study or early clinical studies. These could be modified by other factors such as A-F above. Examples of ECG Monitoring in Late Stage: 1. The thorough QT study results in a negative finding as defined by the E14 criteria* at the therapeutic dose, but the supratherapeutic dose (relative to Phase 3 dose) shows mean QTc effects between 10 and 20 ms. If there is reasonable assurance that the higher dose represents drug exposures that are unlikely to be seen in the patient population, only routine ECG monitoring is recommended in late
--	--	--	--

			phase trials. This approach provides reassurance for safety because patients are unlikely to experience a clinically significant QTc effect. 2. The thorough QT study results in a positive finding as defined by the E14 criteria* at the therapeutic dose, with a mean prolongation <20 ms. For drugs with this magnitude of effect on the QTc interval, intensive monitoring of Phase 3 patients is called for. Intensive ECG monitoring in clinical trials has two main objectives. One objective is to provide protection to patients who might have large worrisome QT intervals >500 ms. A second objective is identifying the frequency of marked QT increases (e.g., prolonged QT >500 ms or increases in QTc >60 ms). Given the limitations of collecting ECGs in late stage trials, the focus of the analysis is on outliers, not on central tendency. Other than descriptive statistics, detailed statistical analysis is not expected. This monitoring is intended to be performed locally, without the involvement of a central core laboratory. The timing of ECG collection should be based on the known properties of the drug. All patients should receive baseline, steady-state, and periodic ECGs during the trial. In addition, ECGs should be collected around Tmax at the first dose and/or around steady state in a subgroup of patients or in dedicated studies. ECG collection at around Tmax is not important for drugs with low fluctuations between peak and trough concentrations. If the drug shows a delayed effect in QT prolongation, then the timing of ECG collection should reflect this delay. 3. The thorough QT study results in a negative finding as defined by the E14 criteria* at the therapeutic dose, but the supratherapeutic
--	--	--	--

			dose shows a mean effect between 10 and 20 ms. If suprathreshold exposure is anticipated at the clinical dose only in a well-characterised subgroup, intensive monitoring as described in Example 2 above could be carried out in this subset of the Phase 3 population. In this case, there should be reasonable assurance that the higher exposure is unlikely to be seen in the general patient population. In contrast, if people in the general patient population (who can not be readily identified in advance) will in some cases achieve this higher exposure, intensive ECG monitoring in the Phase 3 population is expected, as in Example 2. 4. The therapeutic dose results in a mean QTc prolongation of >20 ms. For drugs with large QTc prolongation effects, intensive ECG assessment would be appropriate in all patients in Phase 2/3. Because of the risk of TdP, another important use of ECG monitoring in late phase trials would be to assess any risk mitigation strategies (e.g., electrolyte monitoring, dose reduction strategies). Additional ECG assessment over and above what is recommended earlier in the Q&A might also be called for (e.g., 24 hour ECG recording, telemetry, multiple trough ECGs through steady state). The sponsor is encouraged to discuss these approaches with the relevant regulatory agency(ies) prior to initiation of the Phase 3 program. <i>*A negative study as defined by the ICH E14 criteria is an upper one-sided 95% CI of QTc prolongation effect <10 ms.</i>
--	--	--	--

SECTION 2: S7B QUESTIONS AND ANSWERS

1. INTEGRATED RISK ASSESSMENT

#	Date of Approval	Questions	Answers
1.1	June 2026	ICH S7B applies to new chemical entities for human use. ICH E14 applies to all new drugs having systemic bioavailability. Considering that the spectrum of therapeutic modalities is continuously evolving, can the scope of ICH S7B and ICH E14 be aligned and broadened for the applicability of the principles?	ICH S7B refers to new chemical entities for human use in the scope. In full recognition of the scope of ICH E14, factors that assist in the determination of the most informative nonclinical testing strategy to address the potential for delayed ventricular repolarisation for modalities are provided (see ICH S7B Q&A 1.4). Hence the principles in ICH S7B through its Q&A (date June 2026) can be considered applicable to all modalities. This Guideline should be read in conjunction with ICH S7A, ICH M3, ICH S6, ICH S9 and any future modality-specific Guidelines (e.g., ICH S13).
1.2	June 2026	What is the general strategy for use of nonclinical information as part of an integrated risk assessment for delayed ventricular repolarization and torsade de pointes (TdP) that can inform the design of clinical investigations and interpretation of their results?	The ICH S7B Guideline describes a nonclinical strategy for assessing risk of delayed ventricular repolarisation and QT interval prolongation (Section 2.3). A mechanistic understanding of the development of TdP and the emergence of new types of assays have made it possible to obtain more information to assess TdP risk from nonclinical assays. In addition to supporting the planning and interpretation of First-in-Human clinical studies, nonclinical investigations can also contribute to an integrated risk assessment for TdP in later stages of development when clinical data are available. The following are points to consider when using an integrated nonclinical risk assessment including <i>in vitro</i> data and <i>in vivo</i> QT data in combination with clinical QT data as described in ICH E14 Q&As 5.1 & 6.1. Consideration can be given to alternative assays (e.g., <i>in vitro</i>, <i>ex vivo</i>, or <i>in silico</i>) that are qualified to replace the hERG assay or <i>in vivo</i> QT study (see ICH S7B Q&A 1.5). 1. <i>In vitro</i> hERG or qualified alternative assay: To predict whether the hERG block poses a risk of interfering with ventricular

			repolarisation or TdP, evaluation of the hERG safety margin based on results of a best practice assay (see ICH S7B Q&As 1.3 and 2.1) is recommended. If a qualified alternative assay is used, then see ICH S7B Q&A 1.5. ICH S7A Section 2.6 describes considerations for when human metabolite(s) should be assessed. In these cases, the <i>in vitro</i> hERG assay or qualified alternative assay should evaluate the hERG block liability of the metabolite(s). If an <i>in vitro</i> hERG or qualified alternative assay is not performed, then a scientific justification should be provided (see ICH S7B Q&A 1.4). 2. <i>In vivo</i> QT assessment or qualified alternative assay: The effects on the QTc interval should be assessed at exposures that cover the anticipated high clinical exposure scenario. The adequacy of exposure to any major human metabolites should be determined (see ICH S7A Sections 2.3.3.2 & 2.6, and ICH S7B Q&A 3.5). In addition, if the assay is to be used in situations where a thorough QT study is not feasible as described in ICH E14 Q&A 6.1, then the <i>in vivo</i> QT assessment should have sufficient sensitivity to detect a QTc prolongation effect of a magnitude similar to dedicated clinical QT studies (see ICH S7B Q&A 3.4). This additional consideration (sensitivity similar to dedicated clinical QT studies) does not apply to decision making prior to First-in-Human studies or under ICH E14 Q&A 5.1. A drug with low TdP risk would be expected to have (1) a negative finding in an <i>in vitro</i> hERG assay (see ICH S7B Q&A 1.3 for additional details) or in a qualified alternative assay (see ICH S7B Q&A 1.5 for additional details); and (2) no QTc prolongation in an <i>in vivo</i> QT assessment or qualified alternative assay of sufficient sensitivity conducted at exposures of parent compound and major human metabolites that exceed clinical exposures. A scientific justification
--	--	--	---

			should be provided in case the <i>in vitro</i> assay is not informative or feasible (see ICH S7B Q&A 1.4). When there are factors that can confound or limit the interpretation of the nonclinical studies, such as metabolites and heart rate changes, follow-up studies as described by ICH S7B (Section 2.3.5) can be performed to address these specific issues. If any of the assays included in the integrated nonclinical risk assessment suggest an effect at clinical exposures, the drug has a risk of interfering with ventricular repolarisation. Under this scenario, the drug's TdP risk could be affected by various other factors, such as blocking of additional repolarisation currents (e.g., I_{Ks}), blocking of inward currents (e.g., sodium and L-type calcium currents), effects on the trafficking of ion channel proteins from cytoplasmic sites to the surface membrane, metabolites with ion channel activities, and non-ion channel mediated QT prolongation. Follow-up studies (ICH S7B Section 2.3.5) could be performed to further explore the mechanisms and assess the TdP risk. If applicable, best practice considerations should be followed for assessment of additional ion channel currents (see ICH S7B Q&As 2.1), <i>in vitro</i> cardiomyocyte assays (see ICH S7B Q&As 2.2–2.5), or <i>in vivo</i> studies (see ICH S7B Q&As 3.1–3.6). An appropriately qualified proarrhythmia risk prediction model (see ICH S7B Q&As 4.1–4.2) could be used according to its context of use to assess the possibility of TdP in humans. The assessment of TdP risk using these follow-up studies although optional, can be used together with other relevant nonclinical and clinical information to contribute to the design of subsequent clinical investigations and interpretation of their results.
--	--	--	--

1.3	June 2026	What is the recommended method to compute the hERG safety margin?	A drug's potency for hERG block, usually calculated as half-inhibitory concentration (IC_{50}), can be normalised to the drug's estimated clinically relevant exposures in patients to calculate the safety margin. As more information is obtained during the clinical development, the estimated values of clinical exposures can be refined. When estimating hERG block potency, it is recommended to use standardised procedures and to consider the principles described in S7B Q&A 2.1. The free drug exposure is computed based on the drug's total plasma concentration and the fraction of protein binding. Because of uncertainties in the protein binding measurements, the unbound (free) fraction in plasma should be set to 1% if experimentally determined to be $< 1\%$. This approach has been used in the regulatory assessment for the risk of drug-drug interactions. If protein binding values cannot be accurately assessed (e.g., questionable validation of the bioanalytical method, deviations from best practices), safety margins should be calculated for both steady-state free and total C_{max}. In terms of the exposure to use in the denominator of the safety margin calculation for supporting decision making under ICH E14 Q&As 5.1 or 6.1, it is generally recommended to use the high clinical exposure as defined in ICH E14 Q&A 5.1 (i.e., mean steady state maximum concentration [$C_{max,ss}$] when the maximum therapeutic dose is administered in the presence of the intrinsic or extrinsic factor that has the largest effect on increasing $C_{max,ss}$). To assess whether the hERG block poses a risk of delaying ventricular repolarisation or TdP, the resulting safety margin should be compared to the range of safety margins computed under the same experimental protocol for a series of reference drugs that have known clinical TdP risk and cover diverse electrophysiological properties. Additional pharmacological principles or modeling can be used to justify the use of a certain safety margin threshold (e.g., relationship between hERG block and QTc prolongation from semi-mechanistic
-----	-----------	---	--

			pharmacokinetic/pharmacodynamic models or systems pharmacology models; see Leishman et al. J Pharmacol Toxicol Methods 2020 [https://doi.org/10.1016/j.vascn.2020.106900], for examples). This should be supported by experimental data based on the principles in these Q&As (e.g., same experimental protocol applied to a series of drugs with known clinical TdP risk). Data supporting the safety margin threshold to define a drug as having low TdP risk should be supplied in or appended to the submitted study report. If a recommended hERG margin threshold is published based on principles in these Q&As, a sponsor (or a contract laboratory) seeking to use the same threshold should demonstrate that the inter-laboratory variability of IC₅₀s from a set of calibration drugs under the same experimental protocol does not significantly decrease the sensitivity of the safety margin threshold for detecting drugs that are not low risk for TdP. Appropriate statistical methods should be applied to quantify experimental IC₅₀ variability and calculate uncertainty of the safety margin as confidence/credible intervals.
1.4	June 2026	ICH S7B states that according to current understanding, both an <i>in vitro</i> I_{Kr} (hERG) assay and an <i>in vivo</i> QT assay should be conducted. Considering that the spectrum of therapeutic modalities has changed and that the current understanding of proarrhythmic risk assessment has evolved, what is the framework for identifying the most informative non-clinical testing strategy to address the potential for delayed ventricular repolarisation via direct hERG interaction?	When systemic exposure cannot be ruled out, compounds should generally be tested according to ICH S7A and ICH S7B, including the ICH S7B/E14 Q&As. Considering that the potential for delayed ventricular repolarisation via direct hERG interaction is driven by the physicochemical properties of the modality, the framework for the identification of the most informative nonclinical testing strategy is as follows: For certain modalities, it is acknowledged that the potential to induce delayed ventricular repolarisation via interaction with the hERG channel is not likely. Therefore, no nonclinical studies according to ICH S7B are required, i.e. there is no need for an <i>in vitro</i> hERG assay (see ICH S7B) nor an <i>in vivo</i> QT assessment in line with ICH S7B Q&As 3.1-3.6 for these modalities. Examples include:

			• Monoclonal antibodies (mAb) and large targeted proteins. These include large amino acid (AA) polymers (whether chemically modified or not) that have low membrane permeability and/or are too large to access the common intracellular binding sites on hERG channels even if they were to cross the cell membrane. • AA polymers of any size comprised of naturally occurring amino acids, that the human body generates or encounters through diet, which include the proteinogenic L-AAs and D-AAs. • Certain classes of oligonucleotide-based therapeutics that are considered to have a low likelihood for delayed ventricular repolarisation based on accrued evidence. b. For modalities that are capable of crossing the cell membrane and that can access binding sites within the hERG channel pore (e.g., small molecule), the potential for delayed ventricular repolarisation through direct hERG interaction is likely. In that case, the non-clinical testing strategy will comprise of an <i>in vitro</i> hERG assay (see ICH S7B Q&As 1.2 and 2.1) and an <i>in vivo</i> QT assay preferably integrated into the general toxicology study (see ICH S7B Q&As 3.1-3.6). Where appropriate, consideration can be given to alternative assays (e.g., <i>in vitro</i>, <i>ex vivo</i>, or <i>in silico</i>) that are qualified to replace the hERG assay or the <i>in vivo</i> QT assessment as a core component of the integrated risk assessment (see ICH S7B Q&A 1.5) c. For modalities with unknown potential for delayed ventricular repolarisation through direct hERG interaction, the nonclinical testing strategy is in general comprised of an <i>in vitro</i> hERG assay (see ICH S7B Q&As 1.2 and 2.1) and an <i>in vivo</i> QT assessment that is preferably integrated into the general toxicology study (see ICH S7B Q&As 3.1-3.6). Similar to what has been discussed in ICH S7B Q&A 1.4 part b, where appropriate, consideration can
--	--	--	---

			be given to alternative assays (e.g., <i>in vitro</i>, <i>ex vivo</i> or <i>in silico</i>) that are qualified to replace the hERG assay or <i>in vivo</i> QT assessment as a core component of the integrated risk assessment (see ICH S7B Q&A 1.5). An individualised nonclinical testing strategy should be considered depending on the pharmacodynamic, pharmacokinetic and safety profile of the compound (see ICH S7B). A scientific justification should be provided in case <i>in vitro</i> assays are not informative or feasible. The above considerations can be applicable to novel oligonucleotide-based therapeutics and other novel modalities capable of crossing the cell membrane but without substantial accrued evidence to support a low likelihood of hERG interactions. Some investigational modalities are conjugated molecules (e.g. Antibody-Drug Conjugates), comprised of different moieties. If deconjugation occurs and individual moieties are systemically available, then each moiety should be tested in accordance with its likelihood to cause direct hERG interaction as described under sections a, b and c above. Modalities could be reassigned from “unknown” to “not-likely” or “likely” regarding their potential for delayed ventricular repolarisation through direct hERG interaction based on integrated assessment of physicochemical properties, mechanistic plausibility, and accrued evidence.
1.5	June 2026	In addition to follow up studies, can alternative assays (e.g., <i>in vitro</i> , <i>ex vivo</i> or <i>in silico</i>) be used as a core component of an integrated risk assessment under ICH S7B or clinical scenarios under ICH E14 Q&As 5.1 and 6.1?	Yes, when the alternative assays are qualified for this context of use. The elements outlined in the best practices (see ICH S7B Q&As 2.2-2.5) and general principles (see ICH S7B Q&As 4.1-4.2) should be considered for qualification, where applicable. For example, for <i>in vitro</i> human cardiomyocyte assays used for clinical scenarios under ICH E14 Q&As 5.1 & 6.1, drug-induced changes that inform repolarisation delay should be compared to the reference agents in a manner similar to safety margin comparison described for the hERG assay (see ICH S7B Q&A 1.3).

2. BEST PRACTICE CONSIDERATIONS FOR *IN VITRO* STUDIES

#	Date of Approval	Questions	Answers
2.1	February 2022	What are some “best practice” considerations when evaluating drug potency on affecting cardiac ionic currents using patch clamp method and overexpression cell lines?	As outlined in ICH S7B, the <i>in vitro</i> I_{Kr}/hERG assay plays a critical role in assessing the risk for delayed repolarisation and QT interval prolongation prior to first administration in humans. Nonclinical investigations can also contribute to an integrated risk assessment in later stages of development when clinical QT data are available. The following “best practice” aspects should be considered when sponsors are using I_{Kr}/hERG data to support interpretation of clinical QT data in specific scenarios as described in S7B Q&As 1.2 & 1.3 and ICH E14 Q&As 5.1 & 6.1, and when using Cav1.2 and Nav1.5 to support a proarrhythmia assessment (ICH S7B Q&As 1.2). It is not the intent of these Q&As to make specific recommendations for a sponsor’s screening activities or for all I_{Kr}/hERG assays to support first administration in humans. Several experimental factors are known to influence the potency of drug effects on cardiac ionic currents. These include the voltage protocols used to evoke specific ionic currents, experimental conditions (such as recording temperature, composition of solutions, manual vs. automated assay systems), data acceptance criteria, and data analysis methods employed. Some recommended best practices are therefore provided to enhance reproducibility of <i>in vitro</i> results and the translation to clinical findings. These recommendations are generalisable to voltage clamp experiments characterising potency of drug inhibition (or potentiation) of cardiac currents. 1. Recording temperature: The effects of some drugs are temperature-sensitive and there is currently no method to predict which molecules exhibit temperature-dependent effects or the magnitude of these

			effects. Thus, patch clamp experiments on cells overexpressing cardiac ion channels, including hERG, $\text{Ca}_v1.2$, and $\text{Na}_v1.5$, should be performed at near physiological temperature (35–37 °C). 2. Voltage protocol: The voltage protocols used to evoke ionic currents should approximate the appropriate elements of a ventricular action potential and be repeated at frequencies that are sufficient to minimise the possibility of missing the effects of a test drug at physiologically relevant heart rates. For hERG, a stimulation frequency of 0.2 – 1 Hz is recommended. For $\text{Ca}_v1.2$ and $\text{Na}_v1.5$ currents, a stimulation frequency of 0.2 Hz is recommended. The voltage protocol should include steps that enable monitoring of cell health and consistent electrophysiological recordings throughout the experiment (i.e., estimation of input and series resistance across time). If high seal resistance is achieved, holding current and input resistance (i.e., measures of passive membrane properties at rest) can be used as indicators of cell health and experimental stability. After application of the test drug and if recording quality remains acceptable, a saturating concentration of a selective blocker should be applied to cells to determine residual background current in individual cells. If prominent, background current should be factored into potency determinations. 3. Recording quality: Seal resistance should be high enough so that the leak conductance at all voltages specified by the voltage protocol and series resistance do not compromise voltage control. The extent of series resistance compensation applied to optimise voltage control should be noted. Stability of the ionic current should be demonstrated with baseline recordings (prior to drug application) of sufficient duration to characterise drug-independent changes (such as current run-down). The time course of drug effects should be monitored until steady state effect is obtained, and each cell can be exposed to one or
--	--	--	---

			more drug concentrations as long as cell health and recording quality remain stable. 4. Primary endpoint measures: The primary derived endpoints are inhibitory concentration such as the IC_{50} value (reported in both micromolar and ng/mL units) and Hill coefficient. If 50% current inhibition could not be achieved, a justification of the highest concentration tested should be provided together with the relation of this concentration to therapeutic free and total drug levels. Where necessary, to isolate the current-of-interest, the background current remaining after a high concentration of selective blocker application should be subtracted. If current inhibition with a selective blocker could not be achieved, leak current can be calculated and subtracted from the current traces. This approach assumes that only the current-of-interest is voltage-dependent, hence evidence and justification should be provided on why it was used. 5. Data summary: Inhibition at each drug concentration for each cell should be provided, along with the mean values of IC_{50} and Hill coefficient (and appropriate measures of data variability). To demonstrate recording quality, the study report should also contain time-course plots of current amplitude, input resistance, holding current for individual cells in control condition followed by drug application, and drug equilibration. If time-dependent changes such as current run-up or run-down in baseline condition were corrected for drug inhibition estimation, the correction method applied should be described. 6. Concentration verification: The concentration of test compound to which the cells were exposed should be verified by applying a validated analytical method to the solution collected from the cell chamber. Both nominal and measured concentrations should be
--	--	--	---

			reported. If the nominal and measured concentrations differ significantly from each other, measured concentrations should be used to construct the concentration-response relationship to estimate IC₅₀ and Hill coefficient. 7. Positive and negative controls: The positive control drug should be one of the “reference drugs” referred to in Q&A 1.3. The positive control drug should be tested using sufficient replicates and two or more concentrations achieving 20-80% block, to demonstrate consistency and reproducibility with the reference drug data. If positive control data fall outside the range of expected values, then the study is inconclusive, and it would not be recommended that the data be used to support the purposes outlined in ICH E14 Q&As 5.1 and 6.1. Vehicle (negative) controls should be included in the experiments. The vehicle should include all non-compound materials in the test article solution such as solubilising agents and preservatives.
2.2	February 2022	What are the relevant endpoints of an informative <i>in vitro</i> human cardiomyocyte repolarisation follow-up study?	As outlined in ICH S7B, follow-up studies (Section 2.3.5) can include <i>in vitro</i> ventricular repolarisation assays. Follow-up studies are not performed for all submissions and are often designed to address specific issues. Since implementation of ICH S7B, new technologies have become available, including assays with human induced pluripotent stem cell derived cardiomyocytes (hiPSC-CMs). S7B Q&As 2.2–2.5 outline best practice considerations when <i>in vitro</i> cardiomyocyte assays are performed as follow-up studies. Drug-induced changes in the intracellular or extracellular action potential waveforms recorded from hiPSC-CM preparations and acutely isolated adult human ventricular myocytes reflect the integrated effect on multiple ionic currents, exchangers, and carriers. Changes in cellular repolarisation recognised as markers of ventricular proarrhythmia include delayed and

			abnormal repolarisation (manifest as early afterdepolarisations, triggered activity or irregular beating) and should be noted. Drug induced effects on cardiomyocyte calcium handling (calcium transients, contractility, or calcium homeostasis) could affect electrophysiological activity. These effects may manifest as abnormal mechanical activity associated with proarrhythmic mechanisms (e.g., prolonged contractions, single or multiple premature contractions linked to triggered electrical activity such as early afterdepolarisations). The assessment of changes in calcium handling is useful for describing the presence or absence of potential proarrhythmic activity. However, such measures are inadequate to fully characterise the electrophysiological effects of a drug on repolarisation.
2.3	February 2022	What elements of the test system need to be considered for an <i>in vitro</i> human cardiomyocyte repolarisation assay?	It is important to describe the biological preparation and technology platform that define baseline electrophysiological characteristics and drug responses. • Biological preparation: The origins of cells studied, and human donor characteristics should be specified. If complex preparations containing hiPSC-CMs are used (e.g., co-cultures, organoids, engineered heart tissues), descriptions of the protocols used in creating these preparations should be provided. For primary human cardiomyocyte preparations, the tissue sources, harvesting, isolation, and enrichment procedures followed should be described. Acceptable morphological and functional inclusion criteria for the preparations as well as electrophysiologic characteristics (including baseline action potential/field potential durations, spontaneous beat frequency and variability [if applicable], resting membrane potential, upstroke characteristics, conduction patterns and/or velocity) should be clearly defined. Estimates of the proportion of preparations fulfilling criteria should be included. • Technology platform: The methodologies used (e.g., transmembrane potential recordings [whole cell patch clamp, sharp electrode, or voltage-sensing dye approaches], extracellular recordings using field

			potentials, visual or impedance-based motion approaches, or calcium-sensing dyes) should be clearly described. The analysis package used for marking and interpreting waveforms should be described, with representative recordings (along with pertinent waveform markings) provided. A description of the plates or chambers used (including presence or absence of flow, substrate composition, recording electrode characteristics) should be provided.
2.4	February 2022	What are important considerations when designing and implementing experimental protocols for <i>in vitro</i> cardiomyocyte repolarisation studies?	Protocols should be designed to address a specific question (e.g., concentration-dependent effects on repolarisation). The rationale of choosing a single- or sequential-dose protocol should be provided. Bath temperature should be stable at physiologic (35–37 °C) temperature. The sampling “window” for data collection should be clearly defined. Deviations from protocols should be clearly described, along with expected consequences. • For spontaneously beating preparations, changes in beating rate influence repolarisation independent of direct drug effects on repolarising currents. Spontaneous beat rates in the absence and presence of drugs should be clearly indicated along with the extent of drug-induced rate changes. The choice and justification of correction formula used when assessing repolarisation effects in such preparations should be provided. Due to limitations of rate correction in spontaneously beating hiPSC-CMs, interpretation of potential repolarisation changes may not be possible when a drug causes a rate change. • For paced preparations, the pacing protocol (pattern and duration) should be described and performed in the presence and absence of test compound. • To demonstrate recording quality, the study report should contain time-course plots of primary endpoints (demonstrating drug equilibration) and other parameters that can be used to infer stability of the preparations and signal recordings.

			- Concentration-dependent repolarisation effects can be derived based on vehicle-corrected and/or baseline subtracted comparisons of drug vs. vehicle treated preparations. For higher throughput multi-well platforms, it is preferable to conduct vehicle and test drug studies on the same plate. The number of replicates (useful for evaluating reproducibility but not inferential statistical testing) should be reported. Power calculations are helpful to establish statistical sensitivity to repolarization endpoints. - It is important to characterise drug exposures during <i>in vitro</i> cardiomyocyte repolarisation studies. For well-based studies, drug exposures could be verified using media sampled from test wells or from “satellite studies” (parallel studies using identical protocols and study conditions conducted without measuring electrophysiologic measurements). With continuous flow systems the sampling of effluent from test chambers is valuable for assessing drug exposures. Exposures should be presented as total drug concentration and free drug concentrations (if plasma protein binding characteristics in the media used is known)
2.5	February 2022	How does one define biological sensitivity of a cardiomyocyte <i>in vitro</i> repolarisation assay?	The electrophysiologic sensitivity of cardiomyocyte preparations should be calibrated with established positive controls to confirm their “fit for purpose” role in defining pharmacological block of cardiac ion channel(s). This is readily accomplished by constructing concentration-response curves with recognised and specific ion current blocking agents. - At minimum, it is important to characterise sensitivity to block of the prominent outward repolarising current I_{Kr}/hERG with specific blocking agents (e.g., E-4031 or dofetilide) over relevant concentration ranges. - Block of the inward L-type calcium current (I_{CaL}) and late sodium current (I_{NaL}) may mitigate delayed repolarisation. Demonstrating sensitivity to specific I_{CaL} (e.g., nifedipine or nisoldipine) and I_{NaL} (e.g., mexiletine or lidocaine) blocking agents is helpful for clarifying

			integrated cellular electrophysiological responses of multi-channel blocking drugs.
--	--	--	---

3. BEST PRACTICE CONSIDERATIONS FOR THE *IN VIVO* QT STUDIES

#	Date of Approval	Questions	Answers
3.1	February 2022	What are best practice considerations for species selection and general design of the (standalone) <i>in vivo</i> QTc assessment?	The most appropriate species should be selected and justified (ICH S7B, Section 3.1.3). It is preferable to use the same animal species in the safety pharmacology and non-rodent toxicity studies to facilitate understanding of the possible relationship between adverse cardiovascular pharmacodynamic effects and structural effects on the heart, and to obtain complementary information on systemic exposure level (toxicokinetics). While it is customary to use conscious freely moving telemeterised animals for the <i>in vivo</i> QTc assessments, the choice of alternative model approaches (e.g., anaesthetised or paced animals) might be justified in certain circumstances to achieve adequate exposures or to overcome specific compound-related challenges (e.g., changes in heart rate, tolerability, or bioavailability limitations in conscious animals).
3.2	February 2022	What should be considered for exposure assessment during the <i>in vivo</i> QTc assessment?	The ICH S7B Guideline states that drug exposures should include and exceed anticipated therapeutic concentrations. If the <i>in vivo</i> QTc data are to be used as part of an integrated risk assessment for situations described in ICH E14 Q&As 5.1 & 6.1, the exposure should cover the anticipated high clinical exposure scenario (see S7B Q&A 1.2). An assessment of exposure in the same animals used for the pharmacodynamic assessment is encouraged. Sampling should take place at relevant timepoints and in a manner that limits interference with

			the pharmacodynamic effects. This could be done by sampling complete pharmacokinetic profiles in the same animals on a separate day after an adequate washout or different animals. At least one pharmacokinetic sample should be obtained during the pharmacodynamic assessment day to demonstrate consistency with the full pharmacokinetic profiles. In certain cases, the analysis of QTc interval together with adequate pharmacokinetic sampling makes it possible to perform dedicated exposure-response modeling similar to concentration-QT analysis for clinical QT studies. This can be helpful when the study should be powered to detect an effect similar to dedicated QT studies in humans (e.g., when using <i>in vivo</i> QTc assessment data as part of an integrated nonclinical and clinical risk assessment as described in ICH E14 Q&A 6.1) as it can reduce the number of animals in accordance with the 3R (reduce/refine/replace) principles. In addition, exposure-response modeling may be helpful in other circumstances when QT prolongation is observed or anticipated based on hERG assay results.
3.3	February 2022	What information is needed to support the choice of heart rate correction method in an <i>in vivo</i> QTc assessment?	Optimally, the sponsor should demonstrate the independence of QTc to RR intervals observed in the study through QTc versus RR plots accompanied by additional information (e.g., number of matched QTc-RR pairs, correlation metric, 95% confidence intervals, p-values). QT-RR interval relationship is also important. Justification of correction factors used for QT measures should be provided when test drugs affect heart rate. In certain cases, individual QT correction based on QT-RR relationship is a preferred method as it is more accurate and sensitive than the general methods such as Bazett, Fridericia, or Van de Water when the test drugs affect heart rate. The main reason for not using correction formulae based on historical data is the fixed rate correction coefficients. Non-rodent species show species-specific and individual differences in their QT-RR relationships.

3.4	February 2022	How should the sensitivity of the assay be evaluated?	The test system used for an <i>in vivo</i> QTc assessment should provide a robust response. Assay sensitivity of relevant functional endpoints should be evaluated and reported to enable data interpretation (in supporting initiating first-in-human studies and/or an integrated nonclinical and clinical integrated risk assessment to be applied under the scenarios in ICH E14 Q&As 5.1 or 6.1) and contextualisation. Demonstration of assay sensitivity can be achieved by defining minimum detectable differences and testing the effects of positive controls. Statistical power calculations could also be provided from historical data from the same laboratory using the identical protocol. If historical positive control data are utilised to justify assay sensitivity or statistical power is calculated from historical control data, then the variance of the present data should be consistent with that seen historically. If study results are to be used to support an integrated nonclinical and clinical risk assessment described in ICH E14 Q&A 6.1, then the study should have sensitivity to detect a QTc prolongation effect of a magnitude similar to dedicated clinical QT studies, taking into consideration inter-species differences in the normal range of values for the QTc interval. The overall sensitivity of the nonclinical assay in comparison to clinical QT studies depends on both the electrocardiographic assessment and the exposure achieved in the <i>in vivo</i> QTc assessment relative to high clinical exposure. This can help reduce the numbers of animals used in accordance with the 3R (reduce/refine/replace) principles. The following hypothetical example is offered for consideration with recognition that the QTc threshold and exposure multiples selected for a particular study should be justified by data obtained in the specific species tested, using recognised reference compounds under conditions consistent with the best practice recommendations set forth in these Q&As. • <i>Hypothetical Example:</i> The minimal detectable difference might be 5 milliseconds if drug exposure in the animal study only covers the high clinical exposure but might be higher if a larger multiple of high
-----	---------------	---	--

			clinical exposure is achieved (e.g., 10 milliseconds if 3X high clinical exposure is achieved; or a higher QTc threshold if an even larger multiple is achieved).
3.5	February 2022	What are the recommended conventions for presenting the pharmacodynamic and pharmacokinetic results of an <i>in vivo</i> QTc assessment?	To facilitate the regulatory review of an <i>in vivo</i> QTc assessment, the following are general recommendations. Pharmacodynamic Content • Summary tables and figures showing absolute mean values, mean percent change from baseline, confidence intervals, and p-values for changes from baseline and vehicle control should be included. • If study results are being used to support ICH E14 Q&A 6.1, then data from positive controls should be included or appended. If historical positive controls are used, then the variance of the present data should be consistent with that seen historically, which can be demonstrated by reporting minimal detectable differences with by time analysis. A statement should be provided that the data for the new drug and the historical data were collected according to the same protocol and statistical analysis plan. If deviations are present, they should be clearly justified. If concentration-QTc modeling is performed, reporting should follow similar principles as for human concentration-QTc modeling (see ICH E14 Q&A 5.1). Pharmacokinetic Content • Tabulations of summary statistics for C_{max} (maximum concentration), AUC (area under the curve), and T_{max} (time to maximum concentration) for the parent drug and metabolites along with plasma concentration vs. time plots (if sufficient samples have been collected to support their calculation) should be provided.

			Individual animal data should be provided.
3.6	June 2026	Can <i>in vivo</i> QTc assessments derived from general toxicology studies be used in an integrated risk assessment in line with ICH S7B?	Yes. To reduce the numbers of animals used in accordance with the 3R (reduce/refine/replace) principles, QTc assessments should be incorporated into general toxicology studies whenever feasible, rather than conducted separately. The <i>in vivo</i> QTc assessments included in the general toxicology studies can be used in an integrated risk assessment (ICH S7B) provided that the rigor of these assessments is similar to that in standalone <i>in vivo</i> QTc assessments. This can be achieved with current or new technologies, provided that the methods have been adequately assessed (ICH M3(R2) Q&A). <i>In vivo</i> QTc assessments should use pharmacologically and statistically sensitive approaches such as telemetry (see ICH S7B Q&As 3.1-3.5). Sampling should take place at relevant timepoints and in a manner that limits interference and optimises cardiovascular evaluation (see ICH S7B Q&A 3.2 for further details).

4. PRINCIPLES FOR PROARRHYTHMIA MODELS

#	Date of Approval	Questions	Answers
---	------------------	-----------	---------

4.1	February 2022	The ICH S7B Guideline (Section 3.1.4) states that directly assessing the proarrhythmic risk of pharmaceuticals that prolong the QT interval would be a logical undertaking and interested parties are encouraged to develop these models and test their usefulness in predicting risk in humans. What are general principles to evaluate whether a proarrhythmic risk prediction model could be used as part of an integrated risk assessment strategy?	Different models, including <i>in silico</i>, <i>in vitro</i>, <i>ex vivo</i> and <i>in vivo</i> models, have the potential to be used as part of an integrated risk assessment strategy to evaluate the proarrhythmic risk of QT-prolonging pharmaceuticals in humans. Use of <i>in vitro</i> and <i>in silico</i> models can also reduce animal use in accordance with the 3R (reduce/refine/replace) principles. Because these models have a common feature of using nonclinical experimental data as input and generating human proarrhythmia risk prediction as output, they can generally be referred to as proarrhythmia risk prediction models. The model input can vary among different models, for example, ion channel pharmacology data as input to <i>in silico</i> models, drug-induced changes in cellular repolarisation and/or arrhythmia events as input to hiPSC-CM models, and drug-induced electrocardiographic changes as input to <i>ex vivo/in vivo</i> models. However, the model output (either discrete risk categories or continuous risk scores) is similar among different models. Such a feature makes it possible to develop generic principles for evaluating the predictivity of proarrhythmia risk prediction models without specifying the type of underlying experimental data used as model input. The following general principles should be applied to all proarrhythmia risk prediction models intended to be used as part of an integrated risk assessment for regulatory purposes. While the main focus of these principles is to evaluate a model's predictivity of TdP risk, they are general enough to guide the development of models predicting different types of proarrhythmia. 1. A defined endpoint consistent with the context of use of the model. Examples of model endpoints (e.g., TdP risk vs QT prolongation risk) can be found in Li et al. Clin Pharmacol Ther 2020 (https://doi.org/10.1002/cpt.1647). 2. A fully disclosed algorithm to translate experimental measurements (model input) to proarrhythmia risk (model output), allowing independent reproduction of the model development
-----	---------------	--	--

			process using the associated training and validation datasets to evaluate the model performance. 3. A defined domain of applicability/scope and limitations of the model. This includes the experimental protocols to generate model input (experimental data capturing pharmacological effects of drug), and the compounds tested should have the same arrhythmic mechanisms covered by the model. 4. A prespecified analysis plan and criteria to assess model predictivity. The analysis plan should include methods to separate the training and validation steps. In the training step, a series of reference compounds is used to adjust the model. In the validation step, another series of reference compounds is used to evaluate the performance of the pre-specified model. The reference compounds used for the training and validation steps should not overlap. 5. A mechanistic interpretation of the model, which describes the relationship between the model inputs and mechanism for the arrhythmia. 6. The uncertainty in the model inputs should be captured and propagated to the model predictions. The experimental variability associated with model input should be quantified using appropriate statistical methods and then translated into probabilities of the predicted risk. After a proarrhythmia risk prediction model has been developed, a process should be followed to evaluate whether the model development complied with these principles. Such a process would support that the model is qualified for the intended context of use as part of an integrated risk assessment for regulatory purposes. Some health authorities have
--	--	--	---

			procedures for the formal qualification of models that allow for a model to be used within the qualified context of use without the regulatory authority needing to reconsider and reconfirm its suitability. Model developers are encouraged to contact a regulatory agency about its specific model qualification procedures. After a model has been qualified, the use of such a model is not limited to the specific facility that submitted the qualification package. However, if another facility intends to use the qualified model, that facility should perform laboratory-specific calibration and validation of the model using a subset of the reference compounds that were originally used to develop the model. An illustrative process of performing laboratory-specific calibration and validation is given in Han et al. J Pharmacol Toxicol Methods 2020 (https://doi.org/10.1016/j.vascn.2020.106890).
4.2	February 2022	How can a sponsor use a model for regulatory submission and what are the limitations?	Sponsors can use results from a qualified proarrhythmia model as one component in the totality of evidence approach to risk assessment under the context of use for which the model was developed and qualified. When a facility intends to use the model to produce data for regulatory submission, a set of control compound(s) should be tested to assess the consistency between the new data and the historical laboratory-specific validation data. The number and type of laboratory-specific calibration and control compounds should be justified. If a proarrhythmia model is included in a regulatory submission, proof of qualification of the model under the guidance of the general principles in this Q&A should be provided in an appendix to the study report. Supportive documentation could include published papers, if the included validation dataset is described in sufficient detail to allow an independent assessment. Importantly, the general principles for model qualification set forth in this Q&A only support the use of a proarrhythmia risk prediction model as part of an integrated risk assessment that incorporates all relevant nonclinical and clinical information.