

Contains Nonbinding Recommendations

Draft – Not for Implementation

Draft Guidance on Lisdexamfetamine Dimesylate

October 2024

This draft guidance, when finalized, will represent the current thinking of the Food and Drug Administration (FDA, or the Agency) on this topic. It does not establish any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the Office of Generic Drugs.

In general, FDA’s guidance documents do not establish legally enforceable responsibilities. Instead, guidances describe the Agency’s current thinking on a topic and should be viewed only as recommendations, unless specific regulatory or statutory requirements are cited. The use of the word *should* in Agency guidances means that something is suggested or recommended, but not required.

Active Ingredient: Lisdexamfetamine dimesylate

Dosage Form: Tablet, chewable

Route: Oral

Strengths: 10 mg, 20 mg, 30 mg, 40 mg, 50 mg, 60 mg

Recommended Study: One in vivo bioequivalence study with pharmacokinetic endpoints

1. Type of study: Fasting
Design: Single-dose, two-treatment, two-period crossover in vivo
Strength: 60 mg
Subjects: Healthy males and non-pregnant, non-lactating females
Additional comments: The tablets should be chewed, then and swallowed with water.

Analytes to measure: Lisdexamfetamine and its metabolite, dextroamphetamine, in plasma

Submit the metabolite data as supportive evidence of comparable therapeutic outcome. For the metabolite, the following data should be submitted: individual and mean concentrations, individual and mean pharmacokinetic parameters, and geometric means and ratios of means for area under the curve and maximum concentration.

Bioequivalence based on (90% CI): Lisdexamfetamine

Waiver request of in vivo testing: 10 mg, 20 mg, 30 mg, 40 mg, and 50 mg strengths based on (i) an acceptable bioequivalence study on the 60 mg strength (ii) acceptable in vitro dissolution testing of all strengths, and (iii) proportional similarity of the formulations across all strengths

Dissolution test method and sampling times: The dissolution information for this drug product can be found in the FDA's Dissolution Methods database, <http://www.accessdata.fda.gov/scripts/cder/dissolution>. Conduct comparative dissolution testing on 12 dosage units for each of all strengths of the test product and reference listed drug.¹ Specifications will be determined upon review of the abbreviated new drug application.

Document History: Recommended October 2017; Revised October 2024

Unique Agency Identifier: PSG_208510

¹ If the reference listed drug is not available, refer to the most recent version of the FDA guidance for industry on *Referencing Approved Drug Products in ANDA Submissions*.