

Contains Nonbinding Recommendations

Draft – Not for Implementation

Draft Guidance on Paclitaxel

November 2024

This draft guidance, when finalized, will represent the current thinking of the Food and Drug Administration (FDA, or the Agency) on this topic. It does not establish any rights for any person and is not binding on FDA or the public. You can use an alternative approach if it satisfies the requirements of the applicable statutes and regulations. To discuss an alternative approach, contact the Office of Generic Drugs.

In general, FDA’s guidance documents do not establish legally enforceable responsibilities. Instead, guidances describe the Agency’s current thinking on a topic and should be viewed only as recommendations, unless specific regulatory or statutory requirements are cited. The use of the word *should* in Agency guidances means that something is suggested or recommended, but not required.

Active Ingredient:	Paclitaxel
Dosage Form:	Powder
Route:	Intravenous
Strength:	100 mg/vial
Recommended Studies:	One in vivo bioequivalence study with pharmacokinetic endpoints and one in vitro bioequivalence study with supportive characterization studies

One in vivo bioequivalence study with pharmacokinetic endpoints:

1. Type of study: Bioequivalence study with pharmacokinetic endpoints
Design: Single-dose, two-way crossover in vivo
Strength: 100 mg/vial (260 mg/m² dose administered in 30 minutes)
Subjects: Breast cancer patients after failure of combination chemotherapy for metastatic disease or relapse within 6 months of adjuvant chemotherapy
Additional comments:
 - a. Submission of a Bio-IND is required prior to the conduct of a bioequivalence in vivo study for a cytotoxic drug product such as paclitaxel (see 21 CFR 320.31).¹

¹ As per 21 CFR § 314.94(a)(9)(iii), the proposed parenteral drug product should be qualitatively (Q1) and quantitatively (Q2) the same as the corresponding reference listed drug product (RLD). It is recommended that human serum albumin be sourced from a CBER licensed facility and comply with USP standards and other applicable requirements. Prospective applicants are recommended to obtain assurance from OGD that the Test product has the same in vitro characteristics as the Reference Standard (RS) prior to conducting any in vivo bioequivalence study(ies).

- b. The pivotal bioequivalence study should be conducted using test (T) product manufactured on the proposed commercial scale. In addition, prior to conducting the in vivo bioequivalence study, prospective applicants are recommended to obtain assurance from FDA that the proposed T product has the same in vitro characteristics as the reference standard (RS).
- c. If the patient's health status prevents fasting, the sponsor may provide a non-high-fat diet during the proposed study provided that both study periods are conducted under same conditions.
- d. If the patient's health status necessitates a dose reduction or any change in the recommended 260 mg/m² dose administered in 30 minutes, they are to be withdrawn from the study.
- e. Patients must have baseline neutrophil counts ≥ 1500 cells/mm³; patients who experience a severe hypersensitivity reaction to paclitaxel should not be rechallenged with the drug; frequent peripheral blood counts are to be performed; prior therapy should have included an anthracycline unless clinically contraindicated; female patients should be nonpregnant and non-lactating; women of childbearing potential should be advised to avoid becoming pregnant while receiving paclitaxel intravenous powder, and men should be advised not to father a child while receiving paclitaxel intravenous powder.
- f. The use of antiemetic prophylaxis is acceptable provided that the patient receives the same prophylaxis in both periods of the study.

Analytes to measure: Unbound and total paclitaxel in plasma

Bioequivalence based on (90% CI): AUC and C_{max} for unbound and total paclitaxel

One in vitro bioequivalence study:

- 1. Type of study: In vitro particle size and size distribution
 Design: In vitro bioequivalence study on at least three lots of both T and reference products
 Strength: 100 mg/vial
 Additional comments: None

Parameters to measure: D₁₀, D₅₀, and D₉₀

Bioequivalence based on (95% upper confidence bound): Population bioequivalence based on D₅₀ and SPAN [(D₉₀-D₁₀)/D₅₀]

Comparative characterization studies:

Comparative physicochemical characterization of the T and the RS products. The comparative study should be performed on at least three batches each of the T product² and RS, and it should include:

- Particle morphology
- Particle surface potential
- Paclitaxel crystallinity
- Fraction of free (in solution) and particle-bound paclitaxel and albumin in reconstituted suspension
- Nature of bond between paclitaxel and albumin
- Oligomeric status of albumin in both the albumin excipient and the finished drug product
- In vitro release kinetics

Waiver request of in vivo testing: Not applicable

Dissolution test method and sampling times: Applicants are encouraged to explore methods to characterize in vitro release.

Document History: Recommended November 2024

Unique Agency Identifier: PSG_216338

² The manufacturing process for the exhibit batches should be reflective of the manufacturing process to be utilized for commercial batches.